

# Gene expression analysis in the parvalbumin-immunoreactive PV1 nucleus of the mouse lateral hypothalamus

F. Girard,<sup>1</sup> Z. Meszar,<sup>1</sup> C. Marti,<sup>1</sup> F. P. Davis<sup>2</sup> and M. Celio<sup>1</sup>

<sup>1</sup>Division of Anatomy and Program in Neuroscience, Université de Fribourg, Route A. Gockel 1, CH1700 Fribourg, Switzerland

<sup>2</sup>Janelia Farm Research Campus, Howard Hughes Medical Institute, Ashburn, VA, USA

**Keywords:** brain, calcium-binding protein, glutamate, lateral tuberal nucleus

## Abstract

A solitary, elongated cluster of parvalbumin-immunoreactive neurons has been previously observed in the rodent ventrolateral hypothalamus. However, the function of this so-called PV1 nucleus is unknown. In this article, we report the results of an unbiased, broad and in-depth molecular characterization of this small, compact group of neurons. The Allen Brain Atlas database of *in situ* hybridization was screened in order to identify genes expressed in the PV1-nucleus-containing area of the hypothalamus, and those that might be co-expressed with parvalbumin. Although GABA is the principal neurotransmitter in parvalbumin-expressing cells in various other brain areas, we found that PV1 neurons express the vesicular glutamate transporter (VGLut) VGLut2-encoding gene *Slc17a6* and are negative for the glutamic acid decarboxylase 1 (*GAD1*) gene. These cells also express the mRNA for the neuropeptides *Adcyap1* and possibly *Nxph4*, express several types of potassium and sodium channels, are under the control of the neurotransmitter acetylcholine, bear receptors for the glial-derived neurotrophic factor, and produce an extracellular matrix rich in osteopontin. The PV1 nucleus is thus composed of glutamatergic nerve cells, expressing some typical markers of long-axon, projecting neurons (e.g. VGLut2), but also co-expressing genes typical of short-axon GABA neurons (e.g. a variety of potassium channels). Hence, neurons of the PV1 nucleus combine physiological characteristics of interneurons with those of projection neurons.

## Introduction

The lateral hypothalamic area (LHA) is a large and heterogeneous region, containing several distinct nuclear groups. Although the LHA was historically described as a 'feeding center', its connectivity with autonomic, sympathetic and mesolimbic systems indicates that it regulates the physiological responses to a range of stimuli (such as food, drugs or stress) (reviewed in Berthoud & Münzberg, 2011). Interest in the LHA has recently re-emerged with the discovery of the orexin/hypocretin system and its importance in regulating the sleep/wake cycle (reviewed in Sakurai *et al.*, 2010; Leininger, 2011). The complex structural organization of the LHA (dense and complex array of cell groups and fiber pathways) hampers an analysis of the anatomical to functional relationships. Such an analysis would be aided by a better characterization of the neurotransmitter and neuropeptide content of its neurons.

The EF-hand calcium-binding protein parvalbumin (Pvalb) is strongly expressed in various regions of the postnatal rodent brain, including the cortex, hippocampus, thalamus, midbrain and cerebellum (Celio, 1990; Schwaller, 2009). On the basis of  $Ca^{2+}$ -binding and release kinetics, Pvalb acts as a slow calcium buffer, playing important roles in modulating intracellular calcium dynamics in neurons

(Baimbridge *et al.*, 1992; Schwaller, 2009). A solitary cluster of Pvalb-immunoreactive neurons, referred to as the PV1 nucleus, has been observed in the ventrolateral division of the medial forebrain bundle of the rodent hypothalamus (Celio, 1990). These neurons are orientated horizontally, and have a bipolar or multipolar form (Meszar *et al.*, 2011). Unlike Pvalb-immunoreactive neurons in most other regions of the brain, those of the PV1 nucleus lack the GABA-synthesizing enzyme GAD and GABA itself, but are immunoreactive for glutamate (Meszar *et al.*, 2011).

Topographical mapping of the gene expression patterns in the mammalian central nervous system is crucial for an understanding of the functional networks. During the past few years, research in the field of neuroscience has entered a new era with the release of *in situ* hybridization (ISH) data at the genomic scale, such as appear in the Allen Brain Atlas (ABA) for mouse gene expression (Lein *et al.*, 2007). Indeed, such data provide powerful tools allowing a systematic analysis of particular neuroanatomical structures at the gene expression level (D'Souza *et al.*, 2008; Olszewski *et al.*, 2008; Thompson *et al.*, 2008; Dong *et al.*, 2009).

In this context, identifying genes co-expressed with Pvalb may yield further insight into the evolution and function of the PV1 nucleus. The microarray approach has proved to be an efficient method for genome-wide expression profiling, but it is a challenging technique for structures that embrace only a small number of cells, such as the PV1 nucleus. Therefore, we undertook a genome-scale ISH analysis to

Correspondence: Marco Celio, as above.  
E-mail: marco.celio@unifr.ch

Received 9 June 2011, revised 27 September 2011, accepted 12 October 2011

identify genes co-expressed with Pvalb in PV1 neurons. For this purpose, the ABA database was screened to pinpoint potential candidates, and ISH was used to demonstrate co-expression with Pvalb.

## Materials and methods

### Animals

C57BL/6J mice were used in this study. Animals (20–30 g in weight) were anesthetized with pentobarbital (100 mg/kg), and perfused with ice-cold 0.9% NaCl. Brains were then removed, immediately frozen on dry ice and stored at  $-70^{\circ}\text{C}$  until use. The project was approved by the veterinary committee for animal research of the Canton of Fribourg, Switzerland.

### In situ hybridization and immunostaining

Digoxigenin-labeled RNA probes were prepared following recommendations from Roche (*DIG application manual for non radioactive in situ hybridization*, 3rd edn). Total RNA from mouse brain (Zyagen, San Diego, CA, USA) was reverse transcribed using oligodT as primer, and this reaction mixture was used as the template for polymerase chain reactions using primers specific for each of the genes tested, as described in the ABA (<http://www.brain-map.org/>). Forward and reverse primers were flanked by T3 and SP6 RNA polymerase core promoter sequences, respectively (T3, AAT-TAACCCTCACTAAAGGG and SP6, GCGATTTAGGTGAC-ACTATAG). All primers were purchased from Microsynth (Balgach, Switzerland). Antisense and sense RNA probes were prepared from each specific polymerase chain reaction template by *in vitro* transcription with either SP6 or T3 RNA polymerase and digoxigenin-labeled UTP (Roche Applied Science, Switzerland). Sense probes were used as the negative control of the hybridization procedure. The quality and yields of the probes were estimated on agarose gel. For Pvalb RNA probe, polymerase chain reaction was performed directly using mouse Pvalb full-length cDNA (purchased from ImaGenes, Berlin, Germany) as template, using the Pvalb-specific primers described in the ABA.

For *Macaca fascicularis* probes, the following primers were used in reverse transcription–polymerase chain reaction experiments using total RNA from *Macaca* brain (a gift from Dr F. Raulf, Novartis, Basel, Switzerland): Serpini1 (forward, ATGTGATGAAAATTG-CCAAT; reverse, ATTCCTGAGACAGCAGCAGC) and Slc17a7 (forward, ATGGGCCCCACCCTTAGAAC; reverse, CTAAACT-TCGTGAGGGGGTT). Primers were flanked by a T3 (forward) and SP6 (reverse) RNA polymerase core promoter sequence as described for mouse probes. Probes were made exactly as described for mouse.

The 12–14  $\mu\text{m}$  coronal brain cryosections were prepared on Superfrost gold slides (Medite, Nunningen, Switzerland), and stored at  $-70^{\circ}\text{C}$  until use. ISH was performed essentially following the recommendations of Roche. Briefly, sections were fixed in 4% paraformaldehyde in phosphate-buffered saline for 20 min at  $4^{\circ}\text{C}$ , rinsed in phosphate-buffered saline, and incubated for 10 min in 0.1 M triethanolamine containing 0.25% acetic anhydride. Sections were sequentially incubated in phosphate-buffered saline/0.1 M glycine/0.3% Triton-X100 for 5 min, phosphate-buffered saline for 5 min, 0.1 M Tris, pH 7.5/50 mM EDTA containing  $1\ \mu\text{g}/\text{mL}$  proteinase K at  $37^{\circ}\text{C}$  for 3–5 min, and postfixed for 5 min in 4% paraformaldehyde (PFA). Sections were prehybridized for 30 min at  $57^{\circ}\text{C}$  in 50% formamide/5 $\times$  saline sodium citrate buffer (SSC), then hybridized for 16 h at  $57^{\circ}\text{C}$  with the respective probes in buffer

containing 50% formamide/5 $\times$  SSC/10% Dextran sulfate/1 $\times$  Denhardt's solution/2.5 mg/mL *Torula* RNA. Sections were washed at the hybridization temperature in, respectively, 2 $\times$  SSC, 1 $\times$  SSC and 0.2 $\times$  SSC, 20 min each, then incubated for 30 min in buffer 1 (0.1 M Tris, pH 7.5/0.15 M NaCl) containing 0.1% Triton-X100 and 1% Roche blocking reagent. Alkaline-phosphatase-conjugated anti-digoxigenin (Roche Applied Science) was applied for 2 h. Sections were washed three times for 15 min in buffer 1, and 15 min in buffer 2 (0.1 M Tris, pH 9.5/0.1 M NaCl/50 mM  $\text{MgCl}_2$ ). Detection was performed with nitroblue tetrazolium (NBT)-5-bromo-4-chloro-3'-indolylphosphate (BCIP) (Roche Applied Science) in buffer 2, in the presence of 100 mM levamisole.

Immunostaining with rabbit anti-Pvalb (Swant, Marly, Switzerland) on 20  $\mu\text{m}$  coronal sections was performed following the standard protocol.

### Informatics

ALLENMINER (v1.5; Davis & Eddy, 2009) was used to retrieve three-dimensional expression files from the ABA (Lein *et al.*, 2007) at 100  $\mu\text{m}$  resolution (XPR files;  $n = 21\ 741$ ) and smoothed 200  $\mu\text{m}$  resolution (SVA files;  $n = 24\ 858$ ) formats (January 2010). A region of interest encompassing the PV1 nucleus was first defined in the ABA coordinate system by specifying those voxels in a cuboid [coronal  $x = (56, 71)$ , horizontal  $y = (60, 67)$ , sagittal  $z = (42, 48)$ ] that did not overlap with the tuberal nucleus, subthalamic nucleus, or anterior hypothalamic nucleus, as defined by the ABA reference atlas. One layer of edge voxels that bordered the tuberal nucleus, subthalamic nucleus, or anterior hypothalamic nucleus structures was removed to reduce the effect of mis-registration artifacts, which especially affect small regions (Davis & Eddy, 2009), yielding a final PV1 region of interest containing 699 voxels (of 100  $\mu\text{m}$  resolution).

ALLENMINER queries were then performed to identify (i) genes that are locally enriched in the PV1 region of interest, and (ii) genes whose PV1 expression is similar to Pvalb. First, the ABA data (100  $\mu\text{m}^3$  format) were evaluated for enrichment of gene expression in the PV1 region of interest relative to its local neighborhood, defined by a 4-voxel-thick shell whose PV1-bordering layer is removed (run modes `roiseparate_adjacent`, `roi_list`, `calcentropy_xpr_roi_results`). Second, the similarity of each expression dataset (200  $\mu\text{m}$  format) to each Pvalb expression series (coronal,  $n = 2$ ; sagittal,  $n = 19$ ) was quantified by the Pearson's correlation coefficient of expression levels across all voxels in the region of interest where expression was registered for both genes (run mode `expr_sim_query`).

The accuracies of the searches were assessed using a benchmark set of genes ( $n = 21$ ) whose Pvalb co-expression was independently determined by ISH. When thresholded to avoid all non-PV1-expressing benchmark genes ( $n = 4$ ), the enrichment and similarity queries achieve 50 and 60% true positive rates and predict 584 and 326 PV1-expressing genes, respectively. Combining the results predicts 843 genes, at an estimated 81% true positive rate. Requiring detection by both search strategies predicts 67 genes, at an estimated 25% true positive rate.

## Results

### Computational analysis of the parvalbumin-immunoreactive PV1 nucleus of the lateral hypothalamus

The ABA ISH database covers over 21 000 genes expressed in the adult mouse brain (Lein *et al.*, 2007), and is an invaluable tool for mapping gene expression patterns in specific brain areas (D'Souza

TABLE 1. List of the genes selected after screening the ABA for expression in the PV1 region

Gene	Complete name	Gene ontology
Adcy2	Adenylate cyclase 2	Adenylate cyclase activity
Adcyap1	Adenylate cyclase-activating polypeptide 1	Neuropeptide hormone activity
Anxa5	Annexin A5	Calcium-dependent phospholipid binding
Asb13	Ankyrin repeat and SOCS box-containing protein 13	Unknown
Cart	Cocaine- and amphetamine-regulated transcript	Neuropeptide/hormone activity
Cdh13	Cadherin 13	Cell adhesion
Cdh23	Cadherin 23 (otocadherin)	Cell adhesion
Chrm2	Cholinergic receptor, muscarinic 2, cardiac	Muscarinic acetylcholine receptor activity
Chrm3	Cholinergic receptor, muscarinic 3, cardiac	Muscarinic acetylcholine receptor activity
Chrm2	Cholinergic receptor, nicotinic, beta polypeptide 2 (neuronal)	Nicotinic acetylcholine receptor activity
Col11a1	Procollagen, type XI, alpha 1	ECM constituent
Crh	Corticotropin-releasing hormone	Hormone activity
Crhr1	Corticotropin-releasing hormone receptor 1	Corticotropin-releasing factor receptor activity
Crtac1	Cartilage acidic protein 1	ECM constituent
Dab1	Disabled homolog 1	Signal transduction
Fndc5	Fibronectin type III domain-containing 5	Unknown
Gad1	Glutamic acid decarboxylase 1	Glutamate decarboxylase activity
Gfra1	Glial cell line-derived neurotrophic factor family receptor alpha 1	Receptor activity
Grm1	Glutamate receptor, metabotropic 1	Glutamate receptor activity
Hcn2	Hyperpolarization-activated, cyclic nucleotide-gated K <sup>+</sup> 2	Potassium channel activity
Kcna1	Potassium voltage-gated channel, shaker-related subfamily, member 1	Potassium channel activity
Kcna2	Potassium voltage-gated channel, shaker-related subfamily, member 2	Potassium channel activity
Kcnab2	Potassium voltage-gated channel, shaker-related subfamily, beta member 2	Potassium channel activity
Kcnab3	Potassium voltage-gated channel, shaker-related subfamily, beta member 3	Potassium channel activity
Kcnc1	Potassium voltage-gated channel, Shaw-related subfamily, member 1	Potassium channel activity
Kcnc2	Potassium voltage-gated channel, Shaw-related subfamily	Potassium channel activity
Kcnk1	Potassium channel, subfamily K, member 1	Potassium channel activity
Lgi2	Leucine-rich repeat LGI family, member 2	Unknown
Lhx1	LIM homeobox protein 1	Transcription factor
Limk1	LIM-domain-containing, protein kinase	Protein kinase activity
Lynx1	Ly6/neurotoxin 1	Acetylcholine receptor inhibitor activity
Lynx2	Ly6/Plaur domain-containing 1	Acetylcholine receptor inhibitor activity
Mfap5	Microfibrillar-associated protein 5	ECM constituent
Nefh	Neurofilament, heavy polypeptide	Neurofilament cytoskeleton organization
Nefl	Neurofilament, light polypeptide	Neurofilament cytoskeleton organization
Nef3	Neurofilament 3, medium	Neurofilament cytoskeleton organization
Nos1	Nitric oxide synthase 1, neuronal	Nitric oxide synthase activity
Ntng1	Netrin G1	ECM interaction/axonogenesis
Nxph4	Neuroexophilin 4	Receptor binding
Oprm1	Opioid receptor, mu 1	Neuropeptide receptor activity
Pdyn	Prodynorphin	Neuropeptide/hormone activity
Penk1	Preproenkephalin 1	Opioid peptide activity
Pib5pa	Phosphatidylinositol (4,5) bisphosphate 5-phosphatase, A	Phosphatidylinositol-3,4,5-trisphosphate 5-phosphatase activity
Pitx2	Paired-like homeodomain transcription factor 2	Transcription factor
Plcb4	Phospholipase beta 4	Phospholipase activity
Pnk1	Paroxysmal non-kinesinogenic dyskinesia	Hydroxylase activity
Ret	Ret proto-oncogene	Transmembrane receptor protein tyrosine kinase activity
Sall3	Sal-like 3	Transcription factor
Scg2	Secretogranin II	Cytokine activity
Scn1a	Sodium channel, voltage-gated, type I, alpha	Sodium channel activity
Scn4B	Sodium channel, type IV, beta	Sodium channel activity
Scrt1	Scratch homolog 1, zinc finger protein ( <i>Drosophila</i> )	Transcription factor
Sh3gl2	SH3-domain GRB2-like 2	Unknown
Sema3f	Sema domain, immunoglobulin domain, short basic domain, secreted, (semaphorin) 3f	Chemorepellent, receptor activity
Serpini1	Serine (or cysteine) peptidase inhibitor, clade I, member 1	Peptidase inhibitor activity
Slc17a6	Solute carrier family 17, member 6 (VGlut2)	Neurotransmitter (glutamate) transport
Slc17a7	Solute carrier family 17, member 7 (VGlut1)	Neurotransmitter (glutamate) transport
Sncg	Synuclein gamma	Unknown
Spp1	Secreted phosphoprotein 1	Cytokine activity. ECM binding
Stxbp6	Syntaxin-binding protein (amysin)	Vesicle-mediated transport
Tac1	Tachykinin 1	Neuropeptide activity
Tac1r	Tachykinin receptor 1	Neuropeptide receptor activity
Tac2	Tachykinin 2	Neuropeptide activity
Tmem163	Transmembrane protein 163	Unknown
Trh	Thyrotropin-releasing hormone	Hormone activity
Uqcrh	Ubiquinol-cytochrome c reductase hinge protein	Ubiquinol-cytochrome c reductase activity

TABLE 1. (Continued)

Gene	Complete name	Gene ontology
Vamp1	Vesicle-associated membrane protein 1	Vesicle-mediated transport
LOC382102	Gene model 1125	Unknown
C920006C10Rik	RIKEN cDNA C920006C10 gene (EFR3-like)	Unknown
A930038C07Rik	RIKEN cDNA A930038C07 gene	Heparin binding, ECM organization
6330403A02Rik	RIKEN cDNA 6330403A02 gene	Unknown
LOC433022	Plcx2, phosphatidylinositol-specific phospholipase C, X domain-containing 2	Phospholipase activity
2300002D11Rik	RIKEN cDNA 2300002D11 gene (Trnp1)	Unknown
4930544G21Rik	RIKEN cDNA 4930544G21 gene (Sphkap)	Unknown
4930572J05Rik	RIKEN cDNA 493057J05 gene	Unknown
9130213B05Rik	RIKEN cDNA 9130213B05 gene (Parm1)	Unknown

Given are the abbreviated and complete names, and known or proposed function (Gene Ontology) for the candidate genes selected in this study.

*et al.*, 2008; Olszewski *et al.*, 2008; Thompson *et al.*, 2008; Dong *et al.*, 2009). Initially, we screened the ABA database for genes that are regionally enriched in the lateral hypothalamus, using the provided tools: (i) the anatomic search tool focusing on the hypothalamus; (ii) the gene finder tool of the Anatomic Gene Expression Atlas focusing on the lateral hypothalamus; and (iii) the neuroblast tool searching for genes with an expression pattern similar to that of Pvalb. This first round of screening was then followed by an automated ALLENMINER search (Davis & Eddy, 2009) for genes that are either enriched in the PV1 nucleus, or that exhibit a Pvalb-like expression pattern (see Materials and methods). From these various screens, we chose to analyze in more detail a list of 76 candidate genes expressed in the region containing the PV1 nucleus (see below). These can be ontologically classified as follows (see Table 1): ion channel activity: sodium (Scn1a, Scn4b), potassium (Hcn2, Kcna1, Kcna2, Kcnab2, Kcnab3, Kcnc1, Kcnc2, Kcnk1); transporter activity: Slc17a6, Slc17a7, Stxbp6, Vamp1; enzyme activity: Adcy2, Gad1, Limk1, Nos1, Pib5pa, Plcb4, Plcx2, Pnkd, Uqcrh; receptor activity: Chrm2, Chrm3, Chrb2, Crhr1, Gfra1, Grm1, Oprm1, Ret, Sema3f, Tac1r; transcription factor activity: Lhx1, Pitx2, Sall3, Scrt1; neuropeptide/neurohormone activity: Adcyap1, Cart, Nxfp4, Crh, Pdyn, Penk1, Tac1, Tac2, Trh; cytoskeleton structural components: Nefh, Nef3, Nefl; extracellular matrix (ECM)/cell adhesion molecules: Cdh13, Cdh23, Col11a1, Crtac1, Mfap5, Ntgn1, Spp1, A930038C07Rik; receptor inhibitor activity: Lynx1, Lynx2; peptidase inhibitor activity Serpin1; miscellaneous: Anxa5, Dab1, Scg2; and unknown function: Asb13, EFR3like, Fnec5, Lgi2, Parm1, Sh3gl2, Sncg, Sphkap, Trnp1, Tmem163, LOC382102, 4930572J05Rik,

63304003A02Rik. The complete list of 843 candidate genes identified with the ALLENMINER search is available as Supporting Information (Tables S1–S4).

#### Co-expression with parvalbumin in the PV1 nucleus

In the mouse brain, the PV1 nucleus can be revealed by either ISH on Pvalb mRNA, or Pvalb immunostaining (Fig. 1) (Meszar *et al.*, 2011). In the ABA, it lies between coronal levels 68 and 72, forming an elongated, longitudinally orientated structure within the lateral hypothalamus (Fig. 2G) (Meszar *et al.*, 2011). Typically, between 5 and 10 Pvalb-positive cells are visible on each coronal section (Fig. 1). We thus reasoned that ISH on adjacent 12–14- $\mu$ m-thick coronal cryosections would resolve a sufficient number of cells co-expressing Pvalb and the candidate genes. Using the selection criteria: (i) a restricted expression pattern; (ii) a good signal-to-noise ratio of the antisense RNA probe; and (iii) a potential functional importance, 27 of the 76 candidate genes were tested by ISH. Among them, 19 were shown here to be co-expressed with Pvalb in the same neurons, whereas 8 were expressed by intermingling Pvalb-negative cells. The results are presented in Figs 2 and 3, and summarized in Table 2. Figure 2 depicts the co-expression in some PV1 neurons of Pvalb with three candidate genes, Vamp1, Adcyap1 and Slc17a6 (solute carrier family 17, member 6, encoding the glutamate transporter VGlut2). Figure 3 further presents co-expression with Pvalb of several additional genes: Serpin1, Spp1, Pib5pa, Nefl, Asb13, Hcn2, Kcna2, Kcnc1 and Scn4b. The co-expression was not apparent within all of the Pvalb-positive cells. This incomplete co-existence was perhaps not surprising, as we




FIG. 1. Localization of the PV1 nucleus within the LHA in the adult mouse brain. (A) Schematic representation of brain areas at coronal level 71, according to the Allen Reference Atlas. BLA, basolateral amygdalar nucleus; BMA, basomedial amygdalar nucleus; CP, caudoputamen; CTX, cerebral cortex; DMH, dorsomedial hypothalamic nucleus; HY, hypothalamus; int, internal capsule; MEA, medial amygdalar nucleus; opt, optic tract; STN, subthalamic nucleus; TH, thalamus; TU, tuberal nucleus; V3, third ventricle; VMH, ventromedial hypothalamic nucleus; ZI, zona incerta. The red point indicates the location of the PV1 nucleus at this coronal level. (B) Pvalb mRNA expression, and (C) Pvalb immunoreactivity in the PV1 nucleus. The PV1 nucleus is highlighted with a red circle in A and B. White arrows point to cell bodies immunoreactive for Pvalb. Scale bar: 200  $\mu$ m (B), 20  $\mu$ m (C). For interpretation of color references in figure legend, please refer to the Web version of this article.




FIG. 2. Expression of several genes in Pvalb-expressing cells of the PV1 nucleus. A/B, C/D and E/F are adjacent coronal brain sections stained by ISH for Vamp1 (A and A' higher magnification), Adcyap1 (C and C'), Slc17a6 (E and E') and Pvalb (B/B', D/D' and F/F'). Square delineates the PV1 nucleus shown at higher magnification in A'–F'. Adjacent horizontal brain sections stained for Pvalb (G and higher magnification G') and Slc17a6 (H and H'). Scale bar: 200  $\mu\text{m}$  (B), 20  $\mu\text{m}$  (B'). Black arrows point to cells co-expressing Pvalb and the tested genes. CTX, cerebral cortex; MN, mammillary nucleus; OT, optic tract; PV1, PV1 nucleus, highlighted by a dashed line in G; 3V, third ventricle.

used 12–14- $\mu\text{m}$ -thick adjacent sections. Indeed, given the size of the Pvalb-immunoreactive cell bodies (20–30  $\mu\text{m}$ ) (Meszar *et al.*, 2011), not all of the cell bodies were present on both of the adjacent sections, explaining the lack of complete co-expression, and making a quantification analysis impossible with this technique. However, the possibility that some genes are preferentially expressed in certain neuronal subpopulations but not in others cannot be excluded. This point will need to be clarified by immunohistochemistry.

Previous immunohistochemical studies suggested that PV1 neurons were negative for GABA, and positive for glutamate (Meszar *et al.*, 2011). The present study confirms these observations, as these cells express Slc17a6/VGlu2, but not Gad1. The expression of VGlu2 is sufficient to define a glutamatergic phenotype in neurons (Takamori *et al.*, 2000). Hence, our data indicate that Pvalb-expressing cells of the PV1 nucleus are glutamatergic, excitatory neurons. Not surprisingly, Slc17a7, which encodes the other vesicular glutamate transporter, VGlu1, is not expressed in PV1 cells (Table 2). Most glutamatergic neurons are indeed known to express one of the two transporters, and these cell populations are probably functionally different.

The LHA harbors several neuronal subpopulations expressing various neuropeptides, such as the orexigenic peptides galanin, orexin/hypocretin, melanin-concentrating hormone, and the anorexiogenic peptides neurotensin and Cart (for review, see Leininger, 2011). Several genes encoding hypothalamic neuropeptides were pinpointed during our screening process. As shown in Fig. 4, although the neurons expressing these peptides sometimes abutted on, or even intermingled with, the Pvalb-expressing neurons, cells that co-expressed Pvalb and one of the tested genes (Tac1 and Tac2, Penk1, Pdyn, Trh, and Cart) were never revealed. Similarly, Pvalb-positive neurons of the PV1 nucleus were never observed to co-express orexin/hypocretin, galanin or melanin-concentrating hormone (Meszar *et al.*, 2011).

#### Co-expression with parvalbumin in other brain areas

Next we looked at whether these genes expressed in the PV1 nucleus were also present in Pvalb neurons in other regions of the brain. Closer examination at Pvalb-expressing cells in GABAergic neurons of the reticular nucleus of the thalamus and the Pvalb-positive basket cells in

TABLE 2. Summary of co-expression with Pvalb in the hypothalamic PV1 nucleus, thalamic reticular nucleus (RT) and hippocampal dentate gyrus (DG)

Gene	Co-expression with Pvalb in			Possible co-expression with Pvalb in Stratum oriens and molecular layer of the cerebellum	
	PV1 nucleus	DG	RT		
Adcyap1	Yes	NE	NE	<i>NE</i>	<i>NE</i>
Asb13	Yes	nt (probable)	nt (probable)	<i>Yes (few)</i>	<i>NE</i>
Cart	No	NE	NE	<i>NE</i>	<i>NE</i>
Crhr1	Yes	NE	Yes	<i>NE</i>	<i>NE</i>
Gad1	No	Yes	Yes	<i>Yes</i>	<i>Yes</i>
Gfra1	Yes	Yes	Yes	<i>Yes</i>	<i>NE</i>
Hcn2	Yes	nt (probable)	Yes	<i>Yes (few)</i>	<i>NE</i>
Kcna2	Yes	Yes	Yes	<i>Yes</i>	<i>Yes</i>
Kcnc1	Yes	Yes	Yes	<i>Yes</i>	<i>NE</i>
Lynx1	Yes	No	Yes	<i>Yes</i>	<i>Yes</i>
Lynx2	Yes	No	No	<i>NE</i>	<i>NE</i>
Nefh	Yes	nt (probable)	Yes	<i>Yes</i>	<i>Yes (few)</i>
Nefl	Yes	Yes	Yes	<i>Yes (few)</i>	<i>Yes (few)</i>
Nef3	Yes	nt (probable)	Yes	<i>Yes (few)</i>	<i>Yes (few)</i>
Pdyn	No	NE	NE	<i>NE</i>	<i>NE</i>
Penk1	No	NE	NE	<i>NE</i>	<i>NE</i>
Pib5pa	Yes	Yes	Yes	<i>Yes (few)</i>	<i>NE</i>
Scn4B	Yes	NE	Yes	<i>NE</i>	<i>NE</i>
Scrt1	Yes	nt (probable)	Yes	<i>Yes (few)</i>	<i>NE</i>
Serpini1	Yes	Yes	Yes	<i>Yes</i>	<i>Yes</i>
Slc17a6	Yes	No	No	<i>NE</i>	<i>NE</i>
Slc17a7	No	Yes	No	<i>NE</i>	<i>NE</i>
Spp1	Yes	No	Yes	<i>NE</i>	<i>NE</i>
Tac1	No	nt (probable)	NE	<i>Yes (few)</i>	<i>NE</i>
Tac2	No	nt (probable)	NE	<i>NE</i>	<i>NE</i>
Trh	No	NE	nt (probable)	<i>NE</i>	<i>NE</i>
Vamp1	Yes	Yes	Yes	<i>Yes</i>	<i>Yes (few)</i>

Yes/no, experimentally verified by ISH (this study); nt (probable), not tested experimentally, but probably expressed according to the ABA data; NE, not expressed, according to the ABA data and our own experimental observations. For the expression in the hippocampal stratum oriens and the molecular layer of the cerebellum, all data are from the ABA (written in italics).

the dentate gyrus revealed the co-existence of several of the genes found to be expressed in the PV1 nucleus (shown in Fig. 5 for Serpini1, Kcna2 and Pib5pa). As aforementioned for the PV1 nucleus, co-expression was not always apparent within all of the visible Pvalb-expressing cells, and we cannot therefore conclude on the basis of this study that the candidate genes are universally expressed in all Pvalb-positive neurons of the investigated areas. By carefully analyzing the ABA ISH data, we also compared the expression of these genes in the molecular layer of the cerebellum and the hippocampal stratum oriens, both containing mainly Pvalb-positive GABAergic interneurons. Results for the 27 genes are summarized in Table 2. The following conclusions can be made from these observations. The only genes specific for the PV1 nucleus are Slc17a6, Lynx2 and Adcyap1. The genes Crhr1, Scn4b and Spp1 also show a restricted expression pattern, being present in both the glutamatergic PV1 neurons and the GABAergic neurons of the reticular nucleus, but absent from the Pvalb neurons in all of the other areas analyzed. Several genes are expressed in virtually all Pvalb cells analyzed (GABAergic and glutamatergic): Asb13, Gfra1, Hcn2, Kcnc1, Kcna2, Serpini1, Vamp1 and Lynx1 (with the exception of the cerebellum for the first four genes).

#### *Serpini1 and Slc17a7 are expressed in the monkey lateral tuberal nucleus*

On the basis of its topography and neuroanatomy, the lateral tuberal nucleus (LTN) in primates was previously suggested to correspond to the PV1 nucleus in rodents (Gerig & Celio, 2007), although

neurochemical homologies have not yet been demonstrated. Indeed, LTN neurons do not express Pvalb but can be revealed with antibodies to somatostatin (Fig. 6A) (Gerig & Celio, 2007). We prepared *M. fascicularis* antisense RNA probes to two genes: one marker of the rodent PV1 nucleus, Serpini1, and one gene encoding vesicular glutamate transporter VGlut1, Slc17a7 (note that only Slc17a6, not Slc17a7, was found to be expressed in the mouse PV1, but, unfortunately, we were unable to obtain probe for *Macaca* Slc17a6). As presented in Fig. 6, we found a positive expression for both genes in the LTN. Although very preliminary, these observations suggest a possible evolutionary conservation in gene expression between the rodent PV1 and its putative monkey counterpart, the LTN. In particular, the expression of Slc17a7 demonstrates that LTN neurons are also glutamatergic. It will be interesting to analyze the genome-wide expression profile of the monkey LTN, in order to determine which of the PV1-expressed genes identified in this study are conserved in their expression pattern in the monkey.

#### Discussion

Several conclusions can be drawn from this study: Pvalb-expressing neurons of the PV1 nucleus are glutamatergic, excitatory neurons, which express the Adcyap1 and Nxph4 peptides, but lack classical hypothalamic neuropeptides (including orexin/hypocretin), and share several genes in common with Pvalb-immunoreactive inhibitory, GABA neurons, found in other brain areas.

According to several transcriptomic studies using gene arrays, very few of the genes that were found to be expressed in glutamatergic or




FIG. 3. Gene expression in the PV1 hypothalamic nuclei. Higher magnification of the PV1 nucleus in adjacent sections stained by ISH for Pvalb (A', B', C', D', E', F', G', H' and I') and Serpini1 (A), Hcn2 (B), Spp1 (C), Pib5pa (D), Kcna2 (E), Scn4b (F), Nefl (G), Kcnc1 (H) and Asb13 (I). Black arrows point to cells co-expressing Pvalb and the different tested genes. Scale bar: 20  $\mu$ m.




FIG. 4. Various genes encoding neuropeptides/neurohormones located in the far basolateral hypothalamus do not co-express with Pvalb in the PV1 nucleus. Adjacent coronal brain sections stained by ISH for Pvalb (A'–F') and various genes encoding hypothalamic neuropeptides/neurohormones: Penk1 (A), Cart (B), Tac1 (C), Pdyn (D), Trh (E) and Tac2 (F). Dashed circles delineate the position of the PV1 nucleus.




FIG. 5. Most of the genes identified in Pvalb PV1 cells are also expressed with Pvalb in the reticular thalamic nucleus and the dentate gyrus of the hippocampus. Adjacent coronal brain sections stained by ISH for Pvalb (A'-F') and Serpini1 (A and B), Kcna2 (C and D) and Pib5pa (E and F). Pictures were taken at the level of the hippocampal dentate gyrus (A-A', C-C', E-E') and reticular thalamic nucleus (B-B', D-D', F-F'). Black arrows in panels A-A', C-C' and E-E' highlight cells double stained for Pvalb and the tested genes. RT, reticular nucleus of the thalamus; VPL, ventrolateral nucleus of the thalamus; VPM, ventromedial nucleus of the thalamus; st, stria terminalis.




FIG. 6. Conservation of gene expression in the monkey LTN. *Macaca fascicularis* brain cryosections hybridized with digoxigenin-labeled antisense RNA probes to Slc17a7 (B and E) and Serpini1 (C and D). An adjacent section was immunostained with anti-somatostatin to visualize the LTN (A) (marked with dashed line in A-C). (D and E) Higher magnifications of B and C, focusing on cells stained within the LTN.

GABAergic neurons (Sugino *et al.*, 2006; Hardt *et al.*, 2008; Subkhankulova *et al.*, 2010) were revealed from the ABA screen. The only exceptions were Kcnc1 (a marker for GABAergic cells) and Nefh and Adcyap1 (reported as typical for glutamatergic cells). Some of the genes expressed in the PV1 nucleus (listed in Table 1) are known to occur in subclasses of GABAergic neurons. These include: Pitx2 (Skidmore *et al.*, 2008; Waite *et al.*, 2011), Scrt (Marin & Nieto, 2006), Scn1a (Ogiwara *et al.*, 2007; Martin *et al.*, 2010), Gfra1 (Pozas & Ibanez, 2005; Canty *et al.*, 2009), Lynx2 (Dessaud *et al.*, 2006) and

Tmem163 (Burré *et al.*, 2007). Of the tested genes, only Lynx2, Slc17a6 and Adcyap1 were expressed exclusively in the glutamatergic neurons of the PV1 nucleus. These excitatory neurons of the PV1 nucleus thus express many genes in common with inhibitory neurons, but also very specific genes. It would be interesting to isolate Pvalb-positive cells from these different brain areas, and to compare their transcriptome with microarrays.

The PV1 neurons express several ion channels, including the potassium channels Kcna2 (Kv1-2), Kcnc1 (Kv3-1) and Hcn2, and the

sodium channel *Scn4b*, and possibly *Kcna1* (Kv1-1), *Kcnab2*, *Kcnab3*, *Kcnc2* (Kv3-2), *Kcnk1* and *Scn1a* (Nav1-1). Immunohistochemical and electrophysiological studies have reported a close relationship between Pvalb and fast-spiking GABAergic neurons in the cerebral cortex (Celio, 1986; Kawaguchi & Kondo, 2002). Both Kv3.1 and Kv3.2 are found in Pvalb cortical interneurons, and play a crucial role in the generation of the fast-spiking phenotype (Chow *et al.*, 1999). The presence of the same K<sup>+</sup> channels in neurons of the PV1 nucleus suggests that they also exhibit fast-spiking activity.

Neurons of the PV1 nucleus are devoid of several neuropeptides typical of LHA, including *Tac1/2*, *Penk1*, *Pdyn*, *Trh*, *Cart* (this study), and orexin, galanin and melanin-concentrating hormone (Meszar *et al.*, 2011). *Adcyap1* is the only exception, together with *Nxph4*. *Adcyap1*, also known as pituitary *Adcyap*, belongs to the vasoactive intestinal peptide (VIP)/glucagon/growth hormone-releasing factor superfamily. Several functions have been attributed to *Adcyap1*, including the control of food consumption, synaptic plasticity, antinociceptive effects, or hypophysiotrophic activity (Vaudry *et al.*, 2000). Tracing experiments have shown that PV1 neurons are mostly projecting towards the periaqueductal gray and the laterodorsal tegmental nucleus (Celio & Saper, 1999). As *Adcyap1r*, the gene encoding an *Adcyap1* receptor, is expressed in these two regions (ABA data; see also Vaudry *et al.*, 2000), it could be suggested that *Adcyap1*, synthesized from PV1 Pvalb-expressing neurons, is released into the laterodorsal tegmental nucleus and periaqueductal gray where it acts by binding to its receptor. Another neuropeptide used by the PV1 neurons might be *Nxph4*, a member of a family of secreted glycoproteins of which another representative, *Nxph1*, is expressed primarily in interneurons (Petrenko *et al.*, 1996), and binds to neurexin1 (Missler & Südhof, 1998). For *Nxph4*, no receptor has as yet been identified.

*Spp1*/osteopontin is a secreted glycoprotein, which binds different integrin matrix receptors, and exerts several biological activities including calcium binding, chemotactic effects on astrocytes in the central nervous system, cell adhesion, cell signaling and cell proliferation, and regulation of the ECM (Denhardt *et al.*, 2001). Interestingly, it has an expression pattern in the whole adult mouse brain that closely mimics that of Pvalb (ABA data). *Spp1* has to be added to the long list of antigens found in the perineuronal net (Celio & Blumcke, 1994; Celio *et al.*, 1998), a special ECM rich in proteoglycans and glycoproteins (Bonneh-Karkay & Wiley, 2009). The *Serpini1* brain expression pattern is also very similar to that of Pvalb. *Serpini1*/Neuroserpin is an axonally secreted peptidase inhibitor of the serpin family. It is believed to be a critical regulator of extracellular proteolytic events associated with synaptic plasticity and regeneration, through the inhibition of tissue plasminogen activator, a major serine protease of the brain ECM (Krueger *et al.*, 1997; Bonneh-Karkay & Wiley, 2009). The ECM is particularly accentuated around Pvalb neurons (Meszar *et al.*, 2011). Thus, *Spp1* and *Serpini1* might be involved in shaping a peculiar ECM around neurons expressing Pvalb.

*Lynx1* and *Lynx2* belong to a family of prototoxins that bind to and regulate the functions of nicotinic acetylcholine receptors, both *in vivo* and *in vitro* (Ibanez-Tallon *et al.*, 2002; Miwa *et al.*, 2006; Tekiney *et al.*, 2009). *Lynx1/2* are glycosylphosphatidylinositol (GPI)-anchored proteins, which physically associate with their receptors in the somatodendritic membranes of many neurons (Dessaud *et al.*, 2006). Neurons of the PV1 nucleus express both *Lynx1* and *Lynx2* (our ISH data), and possibly *Chrb2* and *Chrm2* (ABA data), and receive Chat-positive terminals (our unpublished data), suggesting that modulation of cholinergic receptors might be important for their functioning.

In conclusion, our study illustrates the power of genome-wide ISH results to generate hypotheses, and the importance of higher resolution sequential ISH to determine gene batteries expressed in small numbers of cells in the brain. We expect this approach to be generally applicable to refining neuroanatomy and gaining functional insight into other small brain regions.

## Supporting Information

Additional supporting information can be found in the online version of this article:

Table S1. Genes found by both ALLENMINER searches: (1) PV1 enrichment and (2) Pvalb-like PV1 expression.

Table S2. ALLENMINER results for genes with Pvalb-like expression in PV1.

Table S3. ALLENMINER results for enriched in PV1.

Table S4. Genes found by either ALLENMINER search: (1) PV1 enrichment or (2) Pvalb-like expression similarity.

Please note: As a service to our authors and readers, this journal provides supporting information supplied by the authors. Such materials are peer-reviewed and may be re-organized for online delivery, but are not copy-edited or typeset by Wiley-Blackwell. Technical support issues arising from supporting information (other than missing files) should be addressed to the authors.

## Acknowledgements

We thank Dr Bruno Muller for advice on the design of ISH experiments, Simone Eichenberger for animal care, and Véronique Piccand for technical help. We also thank Dr Friedrich Raulf for the gift of monkey brain RNA and tissue. This work was supported by a Swiss National Foundation grant (NF 3100A0-113524) and a Novartis Foundation grant.

## Abbreviations

ABA, Allen Brain Atlas; ECM, extracellular matrix; ISH, *in situ* hybridization; LHA, lateral hypothalamic area; LTN, lateral tuberal nucleus; Pvalb, parvalbumin; SSC, saline sodium citrate buffer; VGluT2, vesicular glutamate transporter 2.

## References

- Baimbridge, K., Celio, M. & Rogers, J. (1992) Calcium binding proteins in the nervous system. *Trends Neurosci.*, **15**, 303–308.
- Berthoud, H.R. & Münzberg, H. (2011) The lateral hypothalamus as integrator of metabolic and environmental needs: from electrical self-stimulation to opto-genetics. *Physiol. Behav.*, **104**, 29–39.
- Bonneh-Karkay, D. & Wiley, C. (2009) Brain extracellular matrix in neurodegeneration. *Brain Pathol.*, **19**, 573–585.
- Burré, J., Zimmermann, H. & Volkandt, W. (2007) Identification and characterization of SV31, a novel synaptic vesicle membrane protein and potential transporter. *J. Neurochem.*, **103**, 276–287.
- Canty, A.J., Dietze, J., Harvey, M., Enomoto, H., Milbrandt, J. & Ibanez, C.F. (2009) Regionalized loss of parvalbumin interneurons in the cerebral cortex of mice with deficits in GFRalpha1 signaling. *J. Neurosci.*, **29**, 10695–10705.
- Celio, M. (1986) Parvalbumin in most gamma-aminobutyric acid-containing neurons of the rat cerebral cortex. *Science*, **231**, 995–997.
- Celio, M. (1990) Calbindin 28K and parvalbumin in the rat nervous system. *Neuroscience*, **35**, 375–475.
- Celio, M. & Blumcke, I. (1994) Perineuronal nets – a specialized form of extracellular matrix in the adult nervous system. *Brain Res. Brain Res. Rev.*, **19**, 128–145.
- Celio, M. & Saper, C. (1999) *The Parvalbumin Positive Nucleus of the Lateral Hypothalamus*. Neuroscience 1999 Abstract. Society for Neuroscience, San Diego.

- Celio, M., Spreafico, R., DeBiasi, S. & Vitellaro-Zucarello, L. (1998) Perineuronal nets: past and presents. *Trends Neurosci.*, **21**, 510–515.
- Chow, A., Erisir, A., Farb, C., Nadal, M., Ozaita, A., Lau, D., Welker, E. & Rudy, B. (1999) K<sup>+</sup> channel expression distinguishes subpopulations of parvalbumin and somatostatin containing neocortical interneurons. *J. Neurosci.*, **19**, 9332–9345.
- Davis, F. & Eddy, S. (2009) A tool for identification of genes expressed in patterns of interest using the Allen Brain Atlas. *Bioinformatics*, **25**, 1647–1654.
- Denhardt, D., Noda, M., O'Regan, A., Pavlin, P. & Berman, J. (2001) Osteopontin as a means to cope with environmental insults: regulation of inflammation, tissue remodeling and cell survival. *J. Clin. Invest.*, **107**, 1055–1061.
- Dessaud, E., Salaun, D., Gayet, O., Chabbert, M. & De Lapeyriere, O. (2006) Identification of Lynx2, a novel member of the ly-6/neurotoxin superfamily, expressed in neuronal subpopulations during mouse development. *Mol. Cell. Neurosci.*, **31**, 232–242.
- Dong, H., Swanson, L., Chen, L., Fanselow, M. & Toga, A. (2009) Genomic-anatomic evidence for distinct functional domains in hippocampal field CA1. *Proc. Natl. Acad. Sci. USA*, **106**, 11794–11799.
- D'Souza, C., Chopra, V., Varhol, R., Xie, Y., Bohacec, S., Zhao, Y., Lee, L., Bilenky, M., Portales-Casamar, E., He, A., Wasserman, W., Goldowitz, D., Marra, M., Holt, R., Simpson, E. & Jones, S. (2008) Identification of a set of genes showing regionally enriched expression in the mouse brain. *BMC Neurosci.*, **9**, 66.
- Gerig, A. & Celio, M. (2007) The human lateral tuberal nucleus: immunohistochemical characterization and analogy to the rodent PV1 nucleus. *Brain Res.*, **1139**, 110–116.
- Hardt, O., Scholz, C., Kusters, D., Yanagawa, Y., Pennartz, S., Cremer, H. & Bosio, A. (2008) Gene expression analysis defines differences between region specific GABAergic neurons. *Mol. Cell. Neurosci.*, **39**, 418–428.
- Ibanez-Tallon, I., Miwa, J., Wang, H., Adams, N., Crabtree, G., Sine, S. & Heintz, N. (2002) Novel modulation of neuronal nicotinic acetylcholine receptors by association with the endogenous protoxin Lynx1. *Neuron*, **33**, 893–903.
- Kawaguchi, Y. & Kondo, S. (2002) Parvalbumin, somatostatin and cholecystokinin as chemical markers for specific GABAergic interneuron types in the rat frontal cortex. *J. Neurocytol.*, **31**, 277–287.
- Krueger, S., Ghisu, G., Cinelli, P., Gschwend, T., Osterwalder, T., Wolfer, D. & Sonderegger, P. (1997) Expression of neuroserpin, and inhibitor of tissue plasminogen activator in the developing and adult nervous system of the mouse. *J. Neurosci.*, **17**, 8984–8996.
- Lein, E., Hawrylycz, M., Ao, N., Ayres, M., Bensinger, A., Bernard, A., Boe, A., Boguski, M., Brockway, K., Byrnes, E., Chen, L., Chen, L., Chen, T., Chin, M., Chong, J., Crook, B., Czaplinska, A., Dang, C., Datta, S., Dee, N., Desaki, A., Desta, T., Diep, E., Dolbeare, T., Donelan, M., Dong, H., Dougherty, J., Duncan, B., Ebbert, A., Eichele, G., Estin, L., Faber, C., Facer, B., Fields, R., Fischer, S., Fliiss, T., Frensley, C., Gates, S., Glattfelder, K., Halverson, K., Hart, M., Hohmann, J., Howell, M., Jeung, D., Johnson, R., Karr, P., Kawal, R., Kidney, J., Knapik, R. & Kuan, C., Lake, J., Laramie, A., Larsen, K., Lau, C., Lemon, T., Liang, A., Liu, Y., Luong, L., Michaels, J., Morgan, J., Morgan, R., Mortrud, M., Mosqueda, N., Ng, L., Ng, R., Orta, G., Overly, C., Pak, T., Parry, S., Pathak, S., Pearson, O., Puchalski, R., Riley, Z., Rickett, H., Rowland, S., Royall, J., Ruiz, M., Sarno, N., Schaffnit, K., Shapovalova, N., Svisay, T., Slaughterbeck, C., Smith, S., Smith, K., Smith, B., Sodt, A., Stewart, N., Stumpf, K., Sunkin, S., Sutram, M., Tam, A., Teemer, C., Thaller, C., Thompson, C., Varnam, L., Visel, A., Whitlock, R., Wornoutka, P., Wolkey, C., Wong, V., Wood, M., Yaylaoglu, M., Young, R., Youngstrom, B., Yuan, X., Zhang, B., Zwingman, T. & Jones, A. (2007) Genome wide atlas of gene expression in the adult mouse brain. *Nature*, **445**, 168–176.
- Leininger, G.M. (2011) Lateral thinking about leptin: a review of leptin action via the hypothalamus. *Physiol. Behav.*, **104**, 572–581.
- Marin, F. & Nieto, M. (2006) The expression of Scratch genes in the developing and adult brain. *Dev. Dyn.*, **235**, 2586–2591.
- Martin, M., Dutt, K., Papale, L., Dube, C., Dutton, S., De Haan, G., Shankar, A., Tufik, S., Meisler, M., Baram, T., Goldin, A. & Escayg, A. (2010) Altered function of the SCN1A voltage-gated sodium channel leads to gamma-aminobutyric acid-ergic (GABAergic) interneuron abnormalities. *J. Biol. Chem.*, **285**, 9823–9834.
- Meszar, Z., Girard, F., Saper, C. & Celio, M. (2011) The lateral hypothalamic parvalbumin immunoreactive (PV1) nucleus in rodents. *J. Comp. Neurol.*, doi: 10.1002/cne.22789 [Epub ahead of print].
- Missler, M. & Südhof, T. (1998) Neurexophilins form a conserved family of neuropeptide-like glycoproteins. *J. Neurosci.*, **18**, 3630–3638.
- Miwa, J., Stevens, T., King, S., Caldorone, B., Ibanez-Tallon, I., Xiao, C., Fitzsimmonds, R., Pavlides, C., Lester, H., Picciotto, M. & Heintz, N. (2006) The protoxin Lynx1 acts on nicotinic acetylcholine receptors to balance neuronal activity and survival in vivo. *Neuron*, **51**, 587–600.
- Ogiwara, I., Miyamoto, H., Morita, N., Atapour, N., Mazaki, E., Inoue, I., Takeuchi, T., Itohara, S., Yanagawa, Y., Obata, K., Furuichi, T., Hensch, T. & Yamakawa, K. (2007) Na(v)1.1 localizes to axons of parvalbumin positive inhibitory interneurons: a circuit basis for epileptic seizures in mice carrying an Scn1a mutation. *J. Neurosci.*, **27**, 5903–5914.
- Olshewski, P., Cedernaes, J., Olsson, F., Levine, A. & Schioth, H. (2008) Analysis of the network of feeding neuroregulators using Allen Brain Atlas. *Neurosci. Biobehav. Rev.*, **32**, 945–956.
- Petrenko, A., Ullrich, B., Missler, M., Krasnoperov, V., Rosahl, T. & Südhof, T. (1996) Structure and evolution of neurexophilin. *J. Neurosci.*, **16**, 4360–4369.
- Pozas, E. & Ibanez, C. (2005) GDNF and GFRA1 promote differentiation and tangential migration of GABAergic neurons. *Neuron*, **45**, 701–713.
- Sakurai, T., Mieda, M. & Tsujino, N. (2010) The orexin system: roles in sleep/wake regulation. *Ann. NY Acad. Sci.*, **1200**, 149–161.
- Schwaller, B. (2009) The continuing disappearance of pure Ca<sup>2+</sup> buffers. *Cell. Mol. Life Sci.*, **66**, 275–300.
- Skidmore, J., Cramer, J., Martin, J. & Martin, D. (2008) Cre fate mapping reveals lineage specific defects in neuronal migration with loss of Pitx2 function in the developing hypothalamus and subthalamic nucleus. *Mol. Cell. Neurosci.*, **37**, 696–707.
- Subkhankulova, T., Yano, K., Robinson, H. & Livesey, F. (2010) Grouping and classifying electrophysiologically defined classes of neocortical neurons by single cell, whole genome expression profiling. *Front. Mol. Neurosci.*, **3**, 10.
- Sugino, K., Hempel, C., Miller, M., Hattox, A., Shapiro, P., Wu, C., Huang, Z. & Nelson, S. (2006) Molecular taxonomy of major neuronal classes in the adult mouse forebrain. *Nat. Neurosci.*, **9**, 99–107.
- Takamori, S., Rhee, J., Rosenmund, C. & Jahn, R. (2000) Identification of a vesicular glutamate transporter that defines a glutamatergic phenotype in neurons. *Nature*, **407**, 189–194.
- Tekiney, A., Nong, Y., Miwa, J., Lieberam, I., Ibanez-Tallon, I., Greengard, P. & Heintz, N. (2009) A role for Lynx2 in anxiety-related behavior. *Proc. Natl. Acad. Sci. USA*, **106**, 4477–4482.
- Thompson, C., Pathak, S., Jeromin, A., Ng, L., MacPherson, C., Mortrud, M., Cusick, A., Riley, Z., Sunkin, S., Bernard, A., Puchalski, R., Gage, F., Jones, A., Bajic, V., Hawrylycz, M. & Lein, E. (2008) Genomic anatomy of the hippocampus. *Neuron*, **60**, 1010–1021.
- Vaudry, D., Gonzalez, B., Basille, M., Yon, L., Fournier, A. & Vaudry, H. (2000) Pituitary adenylate cyclase-activating polypeptide and its receptors: from structure to functions. *Pharmacol. Rev.*, **52**, 269–324.
- Waite, M., Skidmore, J., Billi, A., Martin, J. & Martin, D. (2011) GABAergic and glutamatergic identities of developing midbrain Pitx2 neurons. *Dev. Dyn.*, **240**, 333–346.